

## BIOGAS AND METHANE YIELD FROM RYE GRASS

Tomáš Vítěz<sup>1</sup>, Tomáš Koutný<sup>1</sup>, Milan Geršl<sup>1</sup>, Jan Kudělka<sup>1</sup>, Nuttakan Nitayapat<sup>2</sup>, Pavel Ryant<sup>3</sup>, Stanislav Hejduk<sup>4</sup>, Tomáš Lošák<sup>5</sup>, Monika Vítězová<sup>3</sup>, Jan Mareček<sup>1</sup>

<sup>1</sup> Department of Agricultural, Food and Environmental Engineering, Faculty of Agronomy, Mendel University in Brno, Zemědělská 1, 613 00 Brno, Czech Republic

<sup>2</sup> Department of Biotechnolgy, Faculty of Agro Industry, Kasetsart University, 50 Ngamwongwan road, Ladyao, Chatuchak, Bangkok 10900, Thailand

<sup>3</sup> Department of Agrochemistry, Soil Science, Microbiology and Plant Nutrition, Faculty of Agronomy, Mendel University in Brno, Zemědělská 1, 613 00 Brno, Czech Republic

<sup>4</sup> Department of Animal Nutrition and Forage Production, Faculty of Agronomy, Mendel University in Brno, Zemědělská 1, 613 00 Brno, Czech Republic

<sup>5</sup> Department of Environmentalistics and Natural Resources, Mendel University in Brno, Zemědělská 1, 613 00 Brno, Czech Republic

### Abstract

VÍTĚZ TOMÁŠ, KOUTNÝ TOMÁŠ, GERŠL MILAN, KUDĚLKA JAN, NITAYAPAT NUTTAKAN, RYANT PAVEL, HEJDUK STANISLAV, LOŠÁK TOMÁŠ, VÍTĚZOVÁ MONIKA, MAREČEK JAN. 2015. Biogas and Methane Yield from Rye Grass. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 63(1): 143–146.

Biogas production in the Czech Republic has expanded substantially, including marginal regions for maize cultivation. Therefore, there are increasingly sought materials that could partially replace maize silage, as a basic feedstock, while secure both biogas production and its quality.

Two samples of rye grass (*Lolium multiflorum* var. *westerwoldicum*) silage with different solids content 21% and 15% were measured for biogas and methane yield. Rye grass silage with solid content of 15% reached an average specific biogas yield  $0.431 \text{ m}^3 \cdot \text{kg}^{-1}$  of organic dry matter and an average specific methane yield  $0.249 \text{ m}^3 \cdot \text{kg}^{-1}$  of organic dry matter. Rye grass silage with solid content 21% reached an average specific biogas yield  $0.654 \text{ m}^3 \cdot \text{kg}^{-1}$  of organic dry matter and an average specific methane yield  $0.399 \text{ m}^3 \cdot \text{kg}^{-1}$  of organic dry matter.

Keywords: biogas, biogas yield, methane yield, rye grass

### INTRODUCTION

In Western European countries, the growing number of agricultural biogas plants causes an increasing demand for energy crops as a feedstock in digestion processes. Under the agroclimatic conditions of Central Europe, maize (*Zea mays* L.) is currently the most competitive crop for biogas production due to its high yield of methane, easy integration into existing farming systems, and good suitability for storage of biomass (Schittenhelm, 2008). High biogas and methane yield and the easy harvest, silage and storage of maize make this crop very suitable as co-substrate for biogas plants (Weiland, 2006). On the other hand the over

fertilization of the soil, and the consequent harmful environmental effects on water and biodiversity, together with the current increase in the price of maize makes it necessary to look for alternative biomass feedstock for biogas production. The search of new cheap co-substrates, with high biogas and methan yield, are nowadays major issues in order to obtain a more economically feasible process in biogas plants in Europe.

This paper presents results of lab-scale experiments conducted in a batch fermenters to determine biogas yield (BY) and methan yield (MY) and methane content in biogas of first cut of rye grass (*Lolium multiflorum* var. *westerwoldicum*).

## MATERIAL AND METHODS

Sowing date of rye grass (*Lolium multiflorum* var. *westerwoldicum*) was 13<sup>th</sup> of August 2012 in Vatín production area, Czech Republic, altitude 560 m above sea level. First cut of grass forage was on 23<sup>rd</sup> of October 2012. In order to investigate the impact of ensiling and storage on the biogas yield, freshly harvested and wilted forage was pressed in laboratory scale silos. The laboratory silos were stored at the temperature of 23 °C for 90 days; silage preparation was performed in three repetitions. After 90 days samples of silages were investigated for pH, dry matter (DM), organic dry matter (ODM), ash, total nitrogen, fiber and metabolizable energy (ME). The composition of silage is shown in Tab. I.


Biogas production and quality from rye grass silage was measured in batch anaerobic fermentors at the temperature of 38 °C according to German Standard VDI 4630. For each variant (fresh and wilted forage, Tab. I) of rye silage eight batch fermenters of volume 3 dm<sup>3</sup> have been used. Batch

fermenters were filled with 2 dm<sup>3</sup> of inoculum. For inoculation, fermenter content from commercial biogas plant was used. The inoculum for the batch anaerobic fermentation tests was specified by analysing the parameters DM, ODM, pH, organic acids and the electrical conduction. Two from eight batch fermenters with inoculum volume of 2 dm<sup>3</sup> were used without any addition of rye grass silage. Six remaining batch fermenters with 2 dm<sup>3</sup> of inoculum were filled with 100g of rye grass silage. The biogas produced was collected in wet gas meters over a defined period of 25 days and was measured daily. Besides other gas components methane (CH<sub>4</sub>) and carbon dioxide (CO<sub>2</sub>) content were measured during the batch fermentation tests using gas detector Drager X-am® 7000 (Dräger SafetyAG& Co.KGaA, Germany). Biogas production was converted to standard conditions (T<sub>0</sub> = 273 K, p<sub>0</sub> = 101 325 Pa). The volume of biogas and methane produced by a sample was converted to BY and MY, by expressing them as m<sup>3</sup> per kg of organic dry matter (ODM) of the substrate.


I: Composition of rye grass silage after 90 days

Variant	pH [-]	Dry matter [%]	Organic dry matter [%]	In dry matter			
				Ash [%]	Total nitrogen [%]	Fiber content [%]	ME [MJ·kg <sup>-1</sup> ]
wilted forage	3.9	25.1	87.1	13.1	16.5	17.1	17.9
fresh forage	4.1	15.2	86.9	13.4	17.1	15.5	17.9

ME – metabolisable energy



I: Biogas cumulative production from rye grass silages with different dry matter content



2: Methan cumulative production from rye grass silages with different dry matter content

## RESULTS AND DISCUSSION

Results of batch anaerobic fermentation tests are presented as cumulated biogas yields per kg of ODM and methan yields per kg of ODM after 25 days retention time as a mean value of three replicates. All samples of ensiled rye grass tested show monophasic curves of cumulated biogas production (Fig. 1, Fig. 2).

Determined biogas yield for silage made from rye grass with DM content ranged from 0.608–0.699 Nm<sup>3</sup> biogas per kg of ODM for rye grass silage with DM content of 21% and 0.460–0.401 Nm<sup>3</sup> biogas per kg of ODM for rye grass silage with DM content of 15% respectively. Our results are in correlation with other autors (Amon *et al.*, 2007; Prochnow *et al.*, 2009; Weiland, 2003; Nielsen *et al.*, 2012) who observed similar values for rye grass silage.

Determined methan yields for silage made from rye grass ranged from 0.406–0.402 Nm<sup>3</sup> methan per kg of ODM for rye grass silage with DM content of 21% and 0.263–0.233 Nm<sup>3</sup> methan per kg of ODM for rye grass silage with DM content of 15% respectively. Our results are in correlation with other autors (Amon *et al.*, 2007; Prochnow *et al.*, 2009; Weiland, 2003) who observed similar values for rye grass silage.

Our work indicates that the rye grass silage in two different DM content is supplementary biomass for biogas plants based on traditional inputs (maize silage) and thus enhance the overall biogas production. In order to obtain maximum biogas yield, some parameters have to be considered when selecting a catch crop for biogas production, namely the crop species, the soil characteristics, the weather conditions, harvest time, the dry matter content, and the nitrogen application. All these parameters have direct impact on maximum biogas yield per kg of ODM.

Preliminary calculations on the economy in using catch crops for biogas production revealed that the economy is very dependent on the biomass yield per hectare and of the costs for harvesting, transportation to the biogas plant and biomass storage. In order to be able to analyse such effects, longer term experiments are required. Our findngs are fully in accordance with other authors (Molinuevo-Salces *et al.*, 2013; Wienforth *et al.*, 2010).

### Acknowledgement

This study was financed by the Internal Grant Agency of the Faculty of Agronomy MENDEL in Brno No. 11/2013 Biomass and Bio-Energetic Transformations and by the project "Inovace předmětů vyučovaných v oborech TO, ZS, ARI systém pro řízení poloprovozních bioplynových stanic" IN2140331.

## REFERENCES

- AMON, T., AMON, B., KRYVORUCHKO, V., MACHMULLER, A., HOPFNER-SIXT, K., BODIROZA, V., HRBEK, R., FRIEDEL, J. 2007. Methane production through anaerobic digestion of various energy crops grown in sustainable crop rotations. *Bioresource Technology*, 98(17): 3204–3212.
- MOLINUEVO-SALCES, B., LARSEN, U.S., AHRING, K. B., UELLENDAHL, H. 2013. Biogas production from catch crops: Evaluation of biomass yield and methane potential of catch crops in organic crop rotations. *Biomass and Bioenergy*, 59: 285–292.
- NIELSEN, A. M., FEILBERG, A. 2012. Anaerobic digestion of energy crops in batch. *Biosystems Engineering*, 112: 248–251.
- PROCHNOW, A., HEIERMANN, M., PLÖCHL, M., LINKE, B., IDLER, C., AMON, T., HOBBS, P. J. 2009. Bioenergy from permanent grassland – A review. *Bioresource Technology*, 100: 4931–4944.
- SCHITTENHELM, S. 2008. Chemical composition and methane yield of maize hybrids with contrasting maturity. *European Journal of Agronomy*, 29(2–3): 72–79.
- VDI-GESELLSCHAFT ENERGIETECHNIK/FACHAUSSCHUSS REGENERATIVE ENERGIEN. 2006. VDI 4630. *Fermentation of organic materials, characterisation of the substrate, sampling, collection of material data, fermentation tests*. Berlin: VDI.
- WEILAND, P. 2003. Production and energetic use of biogas from energy crops and wastes in Germany. *Applied Biochemistry and Biotechnology*, 109(1–3): 263–274.
- WEILAND, P. 2006. Biomass digestion in agriculture: a successful pathway for the energy production and waste treatment in Germany. *Engineering in Life Sciences*, 6(3): 302–309.
- WIENFORTH, B., HERRMANN, A., SIELING, K., OHL, S., HARTUNG, E., TAUBE, F., KAGE, H. 2010. Biogas-Expert – grassland methane yield and short-term N efficiency of biogas residues. *Grassland Science in Europe*, 15: 229–231.

Contact information

Tomáš Vítěz: [tomas.vitez@mendelu.cz](mailto:tomas.vitez@mendelu.cz)